



Catalyzed hairpin assembly of magnetic nanoclusters with single nucleotide discrimination

Minero, Gabriel Khose Antonio; Baber, Robert William ; Fock, Jeppe; Hansen, Mikkel Foug

Publication date:
2018

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):

Minero, G. K. A., Baber, R. W., Fock, J., & Hansen, M. F. (2018). *Catalyzed hairpin assembly of magnetic nanoclusters with single nucleotide discrimination*. Abstract from 3rd Functional DNA Nanotechnology Workshop, Rome, Italy.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Catalyzed hairpin assembly of magnetic nanoclusters with single nucleotide discrimination

Gabriel Antonio S. Minero, Robert William Baber, Jeppe Fock, Mikkel Fougth Hansen

DTU Nanotech, B. 345B, Technical University of Denmark, DK-2800 Lyngby, Denmark

We demonstrate detection and single nucleotide discrimination of DNA targets based on hairpin assembly (HA) combined with an optomagnetic (OM) readout detecting the clustering of magnetic nanoparticles (MNPs). The HA assay¹ is based on two initially 'locked' hairpin stems *H1* and *H2* that are grafted onto two separate populations of 100 nm MNPs (Fig. 1a). A fully matching DNA target *C1* 'unlocks' the stem of *H1* via toehold strand displacement. The unlocked *H1* probe subsequently unlocks the stem of *H2* and displaces the *C1* target to form a stable *H1-H2* hybrid linking two MNPs. The clustering state of the MNPs is detected using the OM technique, which probes the rotational dynamics of MNPs in response to an applied oscillating magnetic field at frequency f .² The MNPs have a magnetic moment and a linked optical anisotropy and the magnetic field results in a modulation of the intensity of light transmitted through the MNP suspension (Fig. 1a). Depending on the hydrodynamic volume V_h , the MNP response shows a phase lag φ with respect to the field excitation. Clustering increases V_h and a phase lag is observed at lower frequencies.² The response of single MNPs was mainly detected at $f > 40$ Hz, whereas that from MNP clusters was detected at $f < 40$ Hz. As the HA signal, we take the average phase at $f < 40$ Hz after 50 min of HA reaction at 40°C. We studied the dose-response curve and limit of detection (LOD) for MNP hairpin probe densities of 100% and 40% and the specificity of the assay by comparing *C1* and *Mut* targets, where *Mut* had a single base mismatch in the toehold-recognizing part of *H1*. Figs. 1b and c show the average phase lag for $f < 40$ Hz vs. concentration of *C1* and *Mut* targets for the two probe densities. For 100% probe density, we found *C1* and *Mut* LODs to fall within the range of 1-10 nM and 10-100 nM, respectively, and 16-fold discrimination of *C1* from *Mut* at 10 nM. For 40% probe density (Fig. 1c), the *C1* LOD was improved to 0.1 nM, but the specificity was reduced. Less accessible toehold regions of highly 'crowded' *H1* and *H2* give better discrimination of the single base mutation in the DNA target.




Fig. 1: OM-HA detection. (a) Schematic of assay and OM setup. Measurements were performed using plastic chips sandwiched between two heaters at 40°C. (b)-(c) Dose-response analysis of the HA triggered by *C1* as well *Mut* targets at the indicated hairpin densities.

References

- [1] B. Li *et al.* (2011) *Nucleic Acids Res.* **39**, e110.
- [2] A. Mezger *et al.* (2015) *ACS Nano* **9**, 7374-7382.